

Kirjallisuusviitteet – Spinocerebellaariataksia tyyppi 11 (SCA11)

Rosenberg RN (Ed): Autosomal dominant cerebellar phenotypes: The genotype will settle the issue. *Neurology* 40: 1329 – 1331, 1990

Harding AE: Ataxic disorders. Kirjassa: *Neurology in clinical practice*, 337 – 346. Toim. WG Bradley, RB Daroff, GM Fenichel and CD Mardsen. Butterworth-Heinemann, Boston 1991(a)

Harding AE: Cerebellar and spinocerebellar disorders. Kirjassa: *Neurology in clinical practice*, 1603 – 1623. Toim. WG Bradley, RB Daroff, GM Fenichel and CD Mardsen. Butterworth-Heinemann, Boston 1991(b)

Grewal RP et al.: Clinical and genetic analysis of a distinct autosomal dominant spinocerebellar ataxia. *Neurology* 51: 1423 – 1426, 1998

Klockgether T and Evert B: Genes involved in hereditary ataxias. *Trends Neurosci* 21: 413 – 418, 1998

Klockgether T, Ludtke R, Kramer B, Abele M, Burk K, Schöls L, Riess O, Laccone F, Boesch S, Lopes-Cendes I, Brice A, Inzelberg R, Zilber N and Dichgans J: The natural history of degenerative ataxia: a retrospective study in 466 patients. *Brain* 121: 589 – 600, 1998

Moseley ML et al.: Incidence of dominant spinocerebellar and Friedreich triplet repeats among 361 ataxia families. *Neurology* 51: 1666 – 1671, 1998

Subramony SH and Nance M: Diagnosis and Management of the inherited ataxias. *The Neurologist* 4: 327 – 338, 1998

Ashizawa T (Ed): Repeats may not be everything in anticipation. *Neurology* 53: 1164 – 1165, 1999

Worth PF et al: Autosomal dominant cerebellar ataxia type III: linkage in a large British family to a 7,6-cM region on chromosome 15q14-21.3. *Am J Hum Genet* 65: 420-426, 1999 **(SCA11)**

Tan EK and Ashizawa T: Genetic testing in spinocerebellar ataxias: defining a clinical role. *Arch Neurol* 58: 191 – 195, 2001 – Rev

Margolis RL: The Spinocerebellar Ataxias: Order Emerges from Chaos. *Curr Neurol and Neurosci Rep* 2: 477 – 456, 2002