

## SCA17 – kirjallisuusviiteet

Koide R et al., A neurological disease caused by an expanded CAG trinucleotide repeat in the TATA-binding protein gene: a new polyglutamine disease? *Hum Mol Genet* 8: 2047 – 2053, 1999

Fujugasaki H et al., CAG-repeat expansion in the TATA box-binding protein gene causes autosomal dominant cerebellar ataxia. *Brain* 124: 1939 – 1947, 2001

Nakamura K et al., SCA17, a novel autosomal dominant cerebellar ataxia caused by an expanded polyglutamine in TATA-binding protein. *Hum Mol Genet* 10: 1441 – 1448, 2001

Zühlke C et al., Different types of repeat expansion in the TATA-binding protein gene are associated with a new form of inherited ataxia. *Eur Hum Genet* 9: 160 – 164, 2001

Maryama H et al., Difference in disease-free survival curve and regional distribution according to subtype of spinocerebellar ataxia: study of 1,286 Japanese patients. *Am J Med Genet* 114: 578 – 583, 2002

Silveira I et al., Trinucleotide repeats in 202 families with ataxia: a small expanded (CAG)<sub>n</sub> allele at the SCA17 locus. *Arch Neurol* 59: 623 – 629, 2002

Maltecca F et al., Intergenerational instability and marked anticipation in SCA17. *Neurology* 61: 1441 – 1443, 2003

de Michele G et al., Dementia, ataxia, extrapyramidal features, and epilepsy: phenotype spectrum in two Italian families with spinocerebellar ataxia type 17. *Neurol Sci* 24: 166 – 167, 2003

Rolfs A et al., Clinical features and neuropathology of autosomal dominant spinocerebellar ataxia (SCA17). *Ann Neurol* 54: 367 – 375, 2003

Stevanin G et al., Huntington's disease-like phenotype due to trinucleotide repeat expansion in the TBP and JPH3 genes. *Brain* 126: 1599 – 1603, 2003

Zühlke C et al., Phenotypical variability of expanded alleles in the TATA-binding protein gene. Reduced penetrance in SCA17? *J Neurol* 250: 161 – 163, 2003

Bauer P et al., Trinucleotide repeat expansion in SCA17/TBP in white patients with Huntington's disease-like phenotypes. *J Med Genet* 41: 230 – 232, 2004

Bruni AC et al., Behavioral disorder, dementia, ataxia, and rigidity in a large family with TATA Box-binding protein mutation. *Arch Neurol* 61: 1314 – 1320, 2004

Hagenah JM et al., Focal dystonia as a presenting sign on spinocerebellar ataxia 17. *Mov Disord* 19: 217 – 220, 2004

Masaya O et al., Possible reduced penetrance of expansion of 44 to 47 CAG/CAA repeats in the TATA-binding protein gene in spinocerebellar ataxia type 17. *Arch Neurol* 61: 209 – 212, 2004

Shatunov A et al., Small de novo duplication in the repeat region of the TATA-box-binding protein gene manifest with a phenotype similar to variant Creutzfeldt-Jacob disease. *Clin Genet* 66: 496 – 501, 2004

Toyoshima Y et al., SCA17 homozygote showing Huntington's disease-like phenotype. *Ann Neurol* 55: 281 – 286, 2004

Chong JA et al., TATA-binding protein (TBP)-like factor (TLF) is a functional regulator of transcription: reciprocal regulation of the neurofibromatosis type 1 and c-fos genes by TLF/TRF2 and TBP. *Mol Cell Biol* 25: 2632 – 2643, 2005

van Roon-Mom WM et al., TATA-binding protein in neurodegenerative disease. *Neurosci* 133: 863 – 872, 2005

Zühlke C et al., Spinocerebellar ataxia type 17: report of a family with reduced penetrance of an unstable Gln<sub>49</sub> TBP allele, haplotype analysis supporting a founder effect for unstable alleles and comparative analysis of SCA17 genotypes. *BMC Medical Genetics* 6:27, 2005

Wu YR et al., Analysis of polyglutamine-coding repeats in the TATA-binding protein in different neurodegenerative diseases. *J Neural Transm* 112: 539 – 546, 2005

Manganelli F et al., Electrophysiologic characterization in spinocerebellar ataxia 17. *Neurology* 66: 932 – 934, 2006