

SCA3 – kirjallisuusviitteet

Nakano KK et al: Machado disease: a hereditary ataxia in Portuguese immigrants to Massachusetts. *Neurology* 22: 49 – 55, 1972

Burt T et al., A dominant hereditary ataxia resembling Machado-Joseph disease in Arnhem Land, Australia. *Neurology* 43: 1750 – 1752, 1993

Takiyama Y et al., The gene for Machado-Joseph disease maps to human chromosome 14q. *Nat Genet* 4: 300 – 304, 1993

George-Hyslop PS et al., Machado-Joseph disease in pedigrees of Azorean descent is linked to chromosome 14. *Am J Hum Genet* 55: 120 – 125, 1994

Kawaguchi Y et al., CAG expansions in a novel gene for Machado-Joseph disease at chromosome 14 q 32.1. *Nat Genet* 8: 213 – 215, 1994

Maciel P et al., Correlation between CAG repeat length and clinical features in Machado-Joseph disease. *Am J Hum Genet* 57: 54 – 61, 1995

Schöls L et al., Trinucleotide expansion within the MJD1 gene presents clinically as spinocerebellar ataxia and occurs most frequently in German SCA patients. *Hum Mol Genet* 4: 1001 – 1005, 1995

Stevanin J et al., Linkage disequilibrium at the Machado-Joseph disease/spinal cerebellar ataxia 3 locus: evidence for a common founder effect in French and Portuguese-Brazilian families as well as a second ancestral Portuguese-Azorean mutation. *Am J Hum Genet* 57: 1247 – 1250, 1995

Twist EC et al., Machado-Joseph disease maps to the same region of chromosome 14 as the spinocerebellar ataxia type 3 locus. *J Med Genet* 32: 25 – 31, 1995

Bürk K et al., Autosomal dominant cerebellar ataxia type I. Clinical features and MRI in families with SCA1, SCA2 and SCA3. *Brain* 119: 1497 – 1505, 1996

Dürr A et al., Spinocerebellar ataxia 3 and Machado-Joseph disease: clinical, molecular, and neuropathological features. *Ann Neurol* 39: 490 – 499, 1996

Higgins JJ et al., Mutations in American families with spinocerebellar ataxia (SCA) type3: SCA3 is allelic to Machado-Joseph disease. *Neurology* 46: 208 – 213, 1996

Klockgether T et al., Repeat length and disease progression in spinocerebellar ataxia type 3. *Lancet* 348: 830, 1996

Maruff P et al., Cognitive deficits in Machado-Joseph disease. *Ann Neurol* 40: 421 – 427, 1996

Kaakkola S ja Rinne R: Ataksiat ja niiden erotusdiagnostiikka (katsaus). *Duodecim* 113: 1773 – 1782, 1997

Grewal RP et al., Clinical and genetic analysis of a distinct autosomal dominant spinocerebellar ataxia. *Neurology* 51: 1423 – 1426, 1998

Klockgether T et al., The natural history of degenerative ataxia: a retrospective study in 466 patients. *Brain* 121: 589 – 600, 1998

Lima M et al., Causes of death in Machado-Joseph disease. A case-control Study in the Azores (Portugal). *Arch Neurol* 55: 1341 – 1344, 1998

Moseley ML et al., Incidence of dominant spinocerebellar and Friedreich triplet repeats among 361 ataxia families. *Neurology* 51: 1666 – 1671, 1998

Murata Y et al., Characteristic magnetic resonance imaging findings in Machado-Joseph disease. *Arch Neurol* 55: 33 – 37, 1998

Onodera O et al., Progressive atrophy of cerebellum and brainstem as a function of age and the size of the expanded CAG repeats in the MJD1 gene in Machado-Joseph disease. *Ann Neurol* 43: 288 – 296, 1998

Schmidt T et al., An isoform of ataxin-3-accumulates in the nucleus of neuronal cells in affected brain regions of SCA3 patients. *Brain Pathol* 8: 669 – 679, 1998

Schöls L et al., Sleep disturbance in spinocerebellar ataxias. Is the SCA3 mutation a cause of restless legs syndrome? *Neurology* 51: 1603 – 1607, 1998

Warrick JM et al., Suppression of a polyglutamine-mediated neurodegeneration in Dorsophila by the molecular chaperone HSP70. *Nature Genetics* 23: 425 – 428, 1999

Schöls L et al., Genetic background of apparently idiopathic sporadic cerebellar ataxia. *Hum Genet* 107: 132 – 137, 2000

Storey E et al., Frequency of spinocerebellar ataxia types 1, 2, 3, 6, and 7 in Australian patients with spinocerebellar ataxia. *Am J Med Genet* 95: 351-357, 2000

Tang B et al.: Frequency of SCA1, SCA2, SCA3/MJD, SCA6, SCA7, and DRPLA CAG trinucleotide repeat expansions in patients with hereditary spinocerebellar ataxia from Chinese kindreds. *Arch Neurol* 57: 540 – 544, 2000

Gwinn-Hardy K et al., Spinocerebellar ataxia type 3 phenotypically resembling Parkinson disease in a black family. *Arch Neurol* 58: 296 – 299, 2001

Jardin LB et al., Neurologic findings in Machado-Joseph disease. Relation with disease duration, subtypes, and (CAG)_n. *Arch Neurol* 58: 899 – 904, 2001

Maciel P et al., Improvement in the molecular diagnosis of Machado-Joseph disease. *Arch Neurol* 58: 1821 – 1827, 2001

Bryer A et al., The hereditary adult-onset ataxias in South Africa. *J Neurol Sci* 216: 47-54, 2003

Bürk K et al., Cognitive deficits in spinocerebellar ataxia type 1, 2, and 3. *J Neurol* 250: 207 – 211, 2003

Yoshizawa T et al., Magnetic resonance imaging demonstrates differential atrophy of pontine base and tegmentum in Machado-Joseph disease. *J Neurol Sci* 215: 45 – 50, 2003

Brusco A et al., Molecular genetics of hereditary spinocerebellar ataxia. Mutation analysis of spinocerebellar ataxia genes and CAG/CTG repeat expansion detection in 225 Italian families. *Arch Neurol* 61: 727 – 733, 2004

Gu W et al., The shortest expanded allele of the MJD1 gene in the Chinese MJD kindred with autonomic dysfunction. *Eur Neurol* 52: 107 – 111, 2004

Kawai Y et al., Cognitive impairments in Machado-Joseph disease. *Arch Neurol* 61: 1757-1760, 2004

Lu CS et al., The parkinsonian phenotype of spinocerebellar ataxia type 3 in a Taiwanese family. *Parkinsonism Relat Disord* 10: 369 – 373, 2004

Nandagopal R and Moorthy SGK: Dramatic levodopa responsiveness of dystonia in a sporadic case of spinocerebellar ataxia type 3. *Postgrad Med J* 80: 363 – 365, 2004

van de Warrenburg BPC et al., Peripheral nerve involvement in spinocerebellar ataxias. *Arch Neurol* 61: 257 – 261, 2004

Gould VFC: Mouse models of Machado-Joseph Disease and other polyglutamine spinocerebellar ataxias. *Am Soc Exp Neurother* 2: 480 – 483, 2005

Juvonen V et al. The occurrence of dominant spinocerebellar ataxias among 251 Finnish ataxia patients and the role of predisposing large normal alleles in a genetically isolated population. *Acta Neurol Scand* 111: 154 – 156, 2005

Padiath QS et al., Identification of a novel 45 repeat unstable allele associated with a disease phenotype at the MJD1/SCA3 locus. *Am J Med Genet B Neuropsychiatr* 133: 124 – 126, 2005

van de Warrenburg BPC et al., Age at onset variance analysis in spinocerebellar ataxias: a study in a Dutch-Frech cohort. *Ann Neurol* 57: 505 – 512, 2005

Yeh T-H et al., Autonomic dysfunction in Machado-Joseph disease. *Arch Neurol* 62: 630 – 636, 2005