

SCA4 – kirjallisuusviitteet

Gardner K et al., Autosomal dominant spinocerebellar ataxia: clinical description of a distinct hereditary ataxia and genetic localization to chromosome 16 (SCA4) in a Utah Kindred. *Neurology* 44: A361, 1994

Flanigan K et al., Autosomal dominant spinocerebellar ataxia with sensory axonal neuropathy (SCA4): clinical description and genetic localization to chromosome 16q22.1. *Am J Hum Genet* 59: 392 – 399, 1996

Kaakkola S ja Rinne R: Ataksiat ja niiden erotusdiagnostiikka (katsaus). *Duodecim* 113: 1773 – 1782, 1997

Nachmanoff DB et al., Hereditary ataxia with sensory neuronopathy: Biemond's ataxia. *Neurology* 48: 273 – 275, 1997

Fu Y-H et al., Spinocerebellar Ataxia Type 4. Kirjassa: *Handbook of Ataxia Disorders*, ss. 426 - 433. Ed. Klockgether T, Marcel Dekker Inc., New York-Basel, 2000

Nagaoka U et al., A gene on SCA4 locus causes dominantly inherited pure cerebellar ataxia. *Neurology* 54: 1971 – 1975, 2000

Hirano R et al., Fine mapping of 16q-linked autosomal dominant cerebellar ataxia type III in Japanese families. *Neurogenetics* 5: 215 – 221, 2004

Hallenbroich Y et al., Spinocerebellar ataxia type 4. Investigation of 34 candidate genes. *J Neurol* 252: 1472 – 1475, 2005

Hallenbroich Y et al., Spinocerebellar ataxia type 4 (SCA4): Initial pathoanatomical study reveals widespread cerebellar and brainstem degeneration. *J Neural Transm*, 2005

Ishigawa K et al., An autosomal dominant cerebellar ataxia linked to chromosome 16q22.1 is associated with a single-nucleotide substitution in the 5' untranslated region of the gene encoding a protein with spectrin repeat and Rho guanine-nucleotide exchange-factor domains. *Am J Hum Genet* 77: 280 – 296, 2005