

## **SCA5 – kirjallisuusviitteet**

Ranum LPW et al., Spinocerebellar ataxia type 5 in a family descended from the grandparents of President Lincoln maps to chromosome 11. *Nature Genet* 8: 280 – 284, 1994

Kaakkola S ja Rinne R: Ataksiat ja niiden erotusdiagnostiikka (katsaus). *Duodecim* 113: 1773 – 1782, 1997

Zhu S and Gerhard DS: A transcript map of an 800-kp region on human chromosome 11q13, part of the candidate region for SCA5 and BBS1. *Hum Genet* 103(6): 647 – 680, 1998

Ashizawa T (Ed): Repeats may not be everything in anticipation. *Neurology* 53: 1164 – 1165, 1999

Stevanin G et al., Clinical and MRI findings in spinocerebellar ataxia type 5. *Neurology* 53: 1355 – 1357, 1999

Schut LJ et al., Spinocerebellar ataxia Type 5. In: *Kirjassa: Handbook od Ataxia Disorders*, ss. 435 - 445. Ed. Klockgether T, Marcel Dekker Inc., New York-Basel, 2000

Burk K et al., Spinocerebellar ataxia type 5: clinical and molecular genetic features of a German kindred. *Neurology* 62: 327 – 329, 2004

Ikeda Y et al., Spectrin mutations cause spinocerebellar ataxia type 5. *Nature Genet* 38: 184 – 190, 2006

Bauer P et al., Spectrin mutations in spinocerebellar ataxia (SCA). *Bioessays* 28: 785 – 787, 2006