

SCA6 - kirjallisuusviitteet:

Geschwind DH et al., Spinocerebellar ataxia type 6. Frequency of the mutation and genotype-fenotype correlations. Neurology 49: 1247 – 1251, 1997

Ophoff RA et al., Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutations in the Ca²⁺ channel gene CACNL1A4. Cell 87: 543 – 552, 1996

Gomez CM et al, Spinocerebellar ataxia type 6: gaze-evoked and vertical nystagmus, Purkinje cell degeneration, and variable age of onset. Ann Neurol 42: 933 – 950, 1997

Ikeuchi T et al., Spinocerebellar ataxian type 6: CAG repeat expansion in α_{1A} voltage-dependent calcium channel gene and clinical variations in Japanese population. Ann Neurol 42: 879 – 884, 1997

Kaakkola S ja Rinne R: Ataksiat ja niiden erottusdiagnostiikka (katsaus). Duodecim 113: 1773 – 1782, 1997

Zhuchenko O et al., Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the alpha 1A-voltage-dependent calcium channel. Nat Genet 15: 62 – 67, 1997

Yue Q et al., Progressive ataxia due to a missense mutation in a calcium-channel gene. Am J Hum Genet 61: 1078 – 1087, 1997

Klockgether T et al., The natural history of degenerative ataxia: a retrospective study in 466 patients. Brain 121: 589 – 600, 1998

Moseley ML et al., Incidence of dominant spinocerebellar and Friedreich triplet repeats among 361 ataxia families. Neurology 51: 1666 – 1671, 1998

Dichgans M et al., Spinocerebellar ataxia type 6: Evidence for a strong founder effect among German families. Neurology 52: 849 – 851, 1999

Baloh RW and Jen JC: Episodic ataxia type 2 and spinocerebellar ataxia type 6. Kirjassa: Handbook of Ataxia Disorders, 447 - 467. Ed. Klockgether T, Marcel Dekker Inc., New York-Basel, 2000

Schöls L et al., Genetic background of apparently idiopathic sporadic cerebellar ataxia. Hum Genet 107: 132 – 137, 2000

Storey E et al., Frequency of spinocerebellar ataxia types 1, 2, 3, 6, and 7 in Australian patients with spinocerebellar ataxia. Am J Med Genet 95: 351-357, 2000

Takeichi N et al.,: Dissociation of smooth pursuit and vestibulo-ocular reflex cancellation in SCA-6. Neurology 54: 860 – 866, 2000

Tang B et al., Frequency of SCA1, SCA2, SCA3/MJD, SCA6, SCA7, and DRPLA CAG trinucleotide repeat expansions in patients with hereditary spinocerebellar ataxia from Chinese kindreds. Arch Neurol 57: 540 – 544, 2000

Ishikawa K et al., Cytoplasmic and nuclear polyglutamine aggregates in SCA6 Purkinje cells. Neurology 56: 1753 – 1756, 2001

Koh SH et al., Spinocerebellar atxia type 6 and episodic ataxia type 2 in a Korean family. J Korean Med Sci 16: 809 – 813, 2001

Mariotti C et al., Pathogenic effect of an intermediate size SCA-6 allele (CAG)₁₉ in a homozygous patient. Neurology 57: 1502 – 1504, 2001

Piedras-Renteira ES et al., Increased expression of α -1A Ca²⁺ channel currents arising from expanded trinucleotide repeats in spinocerebellar ataxia type 6. J Neurosci 21: 9185 – 9193, 2001

Sinke RJ et al., Clinical and molecular correlations in spinocerebellar ataxia type 6. A study of 24 Dutch families. Arch Neurol 58: 1839 – 1844, 2001

Soong B et al., Metabolic characterization of a spinocerebellar ataxia type 6. Arch Neurol 58: 300 – 304, 2001

Inagaki A et al., Positron emission tomography and magnetic resonance imaging in spinocerebellar ataxia type 2: a study of symptomatic and asymptomatic individuals. Eur J Neurol 12: 725 – 728, 2005

Jiang H et al., Frequency analysis of autosomal dominant spinocerebellar ataxias in mainland Chinese patients and clinical and molecular characterization of spinocerebellar ataxia type 6. Chin Med J 118: 837 – 843, 2005

Juvonen V et al. The occurrence of dominant spinocerebellar ataxias among 251 Finnish ataxia patients and the role of predisposing large normal alleles in a genetically isolated population. Acta Neurol Scand 111: 154 – 1562, 2005

Mascke M et al., Clinical feature profile of spinocerebellar ataxia type 1 – 8 predicts genetically defined subtypes. Mov Disord 20: 1405 – 1412, 2005

van de Warrenburg BPC et al., Age at onset variance analysis in spinocerebellar ataxias: a study in a Dutch-French cohort. Ann Neurol 57: 505 – 512, 2005

Wüllner U et al., Dopamine transporter positron emission tomography in spinocerebellar ataxias type 1, 2, 3 and 6. Arch Neurol 62: 1280 – 1285, 2005