

## **SCA7 – kirjallisuusviitteet**

Benomar A et al., Autosomal-dominant cerebellar ataxia with retinal degeneration (ADCA type II) is genetically different from ADCA type I. Ann Neurol 35: 439 – 444, 1994

Evevoldson TP et al., Autosomal dominant cerebellar ataxia with pigmentary macular dystrophy. A clinical and genetic study of eight families. Brain 117: 445 – 460, 1994

Benomar A et al., The gene for autosomal dominant cerebellar ataxia with pigmentary macular dystrophy maps to chromosome 3p12-p21.1. Nat Genet 10: 84 – 88, 1995

Gouw LG et al., Retinal degeneration characterizes a spinocerebellar ataxia mapping to chromosome 3p. Nat Genet 10: 89 – 93, 1995

David G et al., The gene for autosomal dominant cerebellar ataxia type II is located in a 5cM region in 3p12 – p13: genetic and physical mapping of the SCA7 locus. Am J Hum Genet 59: 1328 – 1336, 1996

Lindblad K et al., An expanded CAG repeat sequence in spinocerebellar ataxia type 7. Genome Res 6: 965 – 971, 1996

David G et al., Cloning of the SCA7 gene reveals a highly unstable CAG repeat expansion. Nat Genet 17: 65 – 70, 1997

Kaakkola S ja Rinne R: Ataksiat ja niiden erotusdiagnostiikka (katsaus). Duodecim 113: 1773 – 1782, 1997

Benton CS et al., Molecular and clinical studies in SCA-7 define a broad clinical spectrum and the infantile phenotype. Neurology 51: 1081 – 1086, 1998

David G et al., Molecular and clinical correlations in autosomal dominant cerebellar ataxia with progressive macular dystrophy (SCA7). Hum Mol Genet 7: 165 – 170, 1998

Gouw LG et al., Analysis of the dynamic mutation in the SCA7 gene shows marked parental effects on CAG repeat transmission. Hum Mol Genet 7: 525 – 532, 1998

Johansson J et al., Expanded CAG repeat in Swedish spinocerebellar ataxia type 7 (SCA7) patients: effect of CAG repeat length on the clinical manifestation. Hum Mol Genet 7: 171 – 176, 1998

Holmberg M et al., Spinocerebellar ataxia type 7 (SCA7): a neurodegenerative disorder with neuronal intranuclear inclusions. Hum Mol Genet 7: 913 – 918, 1998

Klockgether T et al., The natural history of degenerative ataxia: a retrospective study in 466 patients. Brain 121: 589 – 600, 1998

Moseley ML et al., Incidence of dominant spinocerebellar and Friedreich triplet repeats among 361 ataxia families. Neurology 51: 1666 – 1671, 1998

Jonasson J et al., Evidence for common spinocerebellar ataxia type 7 (SCA7) founder mutation in Scandinavia. Eur J Hum Genet 8: 918 – 922, 2000

Modi G et al., The clinical and genetic characteristics of spinocerebellar ataxia type 7 (SCA7) in three Black South African Families. Acta Neurol Scand 101: 177 – 182, 2000

Potter NT and Nance MA: Genetic testing for ataxia in North America. Mol Diagn 5: 91 – 99, 2000

Stevanin G et al., Spinocerebellar Ataxia Type 7. Kirjassa: Handbook of Ataxia Disorders, 470 - 486. Ed. Klockgether T, Marcel Dekker Inc., New York-Basel, 2000

Storey E et al., Frequency of spinocerebellar ataxia types 1, 2, 3, 6, and 7 in Australian patients with spinocerebellar ataxia. Am J Med Genet 95: 351 – 357, 2000

Tang B et al., Frequency of SCA1, SCA2, SCA3/MJD, SCA6, SCA7, and DRPLA CAG trinucleotide repeat expansions in patients with hereditary spinocerebellar ataxia from Chinese kindreds. Arch Neurol 57: 540 – 544, 2000

Jardim LB et al., A survey of spinocerebellar ataxia in South Brazil – 66 new cases with Machado-Joseph disease, SCA7, SCA8, or unidentified disease-causing mutations. J Neurol 248: 870 – 876, 2001

Oh AK et al., Slowing of voluntary and involuntary saccades: an early sign in spinocerebellar ataxia type 7. Ann Neurol 49: 801 – 804, 2001

Yvert G et al., SCA7 mouse models show selective stabilization of mutant ataxin-7 and similar cellular responses in different neuronal cell types. Hum Mol Genet 10: 1679 – 1692, 2001

Bang OY et al., Pontine atrophy precedes cerebellar degeneration in spinocerebellar ataxia 7: MRI-based volumetric analysis. J Neurol Neurosurg Psychiatry 75: 1452 – 1456, 2004

Helmlinger D et al., Disease progression despite early loss of polyglutamine protein expression in SCA7 mouse model. J Neurosci 24: 1881 – 1887, 2004

Helmlinger D et al., Hsp70 and Hsp40 Chaperones do not modulate retinal phenotype in SCA7 mice. J Biol Chem 279: 55969 – 55977, 2004

Bowman AB et al., Neuronal dysfunction in a polyglutamine disease model occurs in the absence of ubiquitin-proteasome system impairment and inversely correlates with the degree of nuclear inclusion formation. Hum Mol Genet 14: 679 – 691, 2005

van de Warrenburg BPC et al., Peripheral nerve involvement in spinocerebellar ataxias. Arch Neurol 61: 257 – 261, 2004

Juvonen V et al. The occurrence of dominant spinocerebellar ataxias among 251 Finnish ataxia patients and the role of predisposing large normal alleles in a genetically isolated population. Acta Neurol Scand 111: 154 – 1562, 2005

Mascke M et al., Clinical feature profile of spinocerebellar ataxia type 1 – 8 predicts genetically defined subtypes. Mov Disord 20: 1405 – 1412, 2005

Rub U et al., Spinocerebellar ataxia type 7 (SCA7): first report of a systematic neuropathological study of the brain of a patient with a very short expanded CAG-repeat. Brain Pathol 15: 287 – 295, 2005

van de Warrenburg BPC et al., Age at onset variance analysis in spinocerebellar ataxias: a study in a Dutch-French cohort. Ann Neurol 57: 505 – 512, 2005

Crum BA and Josephs KA: Varied electrophysiologic patterns in spinocerebellar ataxia type 2. Eur J Neurol 13: 104 – 197, 2006