

SCA8 – kirjallisuusviitteet

Koob MD et al., An untranslated CTG expansion causes a novel form of spinocerebellar ataxia (SCA8). *Nature genetics* 21: 379 – 384, 1999

Day JW et al., Spinocerebellar ataxia type 8. Clinical features in a large family. *Neurology* 55: 649 – 657, 2000

Ikeda Y et al., Molecular and clinical analyses of spinocerebellar ataxia type 8 in Japan. *Neurology* 54: 950 – 955, 2000

Juvonen V et al., Clinical and Genetic Findings in Finnish Ataxia Patients with the Spinocerebellar Ataxia 8 Repeat Expansion. *Ann Neurol* 48: 354 – 361, 2000

Moseley ML et al., SCA8 CTG repeat: en masse contractions in sperm and intergenerational sequence changes may play a role in reduced penetrance. *Hum Mol Genet* 9: 2125 – 2130, 2000

Nemes JP et al., The SCA8 transcript is an antisense RNA to a brain-specific transcript encoding a novel actin-binding protein (KLHL1). *Hum Mol Genet* 9: 1543 – 1551, 2000

Silveira I et al., High germinal instability of the (CTG)_n at the SCA8 locus of both expanded and normal alleles. *Am J Hum Genet* 66: 830 – 840, 2000

Stevanin G et al., Are (CTG)_n expansions at the SCA8 locus rare polymorphisms? *Nature Genetics* 24: 213, 2000

Vincent JB et al., An unstable trinucleotide-repeat region on chromosome 13 implicated in spinocerebellar ataxia: a common expansion locus. *Am J Hum Genet* 66: 819 – 829, 2000

Worth PF et al., Large, expanded repeats in SCA8 are not confined to patients with cerebellar ataxia. *Nature Genetics* 24: 214 – 215, 2000

Cellini A et al., Genetic and Clinical Analysis of Spinocerebellar Ataxia Type 8 Repeat Expansion in Italy. *Arch Neurol* 58 : 1856 – 1859, 2001

Sorbido M-J et al., SCA 8 repeat expansions in ataxia: A controversial association. *Neurology* 57: 1310 – 1312, 2001

Tan E-K and Ashizawa T: Genetic testing in spinocerebellar ataxias. Defining a clinical role. *Arch Neurol* 58: 191 – 195, 2001

Cellini A et al., A Family with spinocerebellar ataxia type 8 expansion and vitamin E deficiency ataxia. *Arch Neurol* 59: 1952 – 1953, 2002

Izumi Y et al., SCA8 repeat expansion: Large CTA/CTG repeat alleles are more common in ataxic patients, including those with SCA6. *Am J Hum Genet* 72: 704 – 709, 2003

Mosemiller AK et al., Molecular genetics of spinocerebellar ataxia type 8 (SCA8). *Cytogenet Genome Res* 100: 175 – 183, 2003

Schols L et al., Do CTG expansions at the SCA8 locus cause ataxia? *Ann Neurol* 54: 5 – 7, 2003

Wu YR et al., Genetic testing in spinocerebellar ataxia in Taiwan: expansion of trinucleotide repeats in SCA8 and SCA17 are associated with typical Parkinson's disease. *Clin Genet* 65: 209 – 214, 2004

Zeman A et al., Spinocerebellar ataxia type 8 in Scotland: genetic and clinical features in seven unrelated cases and a review of published reports. J Neurol Neurosurg Psychiatry 73: 459 – 465, 2004

Baba Y et al., Sporadic SCA8 mutation resembling corticobasal degeneration. Parkinson Relat Disord 11: 147 – 150, 2005

Juvonen V et al. The occurrence of dominant spinocerebellar ataxias among 251 Finnish ataxia patients and the role of predisposing large normal alleles in a genetically isolated population. Acta Neurol Scand 111: 154 – 156, 2005

Lilja A et al., Cognitive impairment in spinocerebellar ataxia type 8. J Neurol Sci 237: 31 – 38, 2005